

Calculus I Students,

Test 3 will be April 1 and April 2 (Augusta students) and April 8 and April 9 (Swainsboro/Statesboro students). To be prepared, you should have done or should do the following:

1. Watched all the videos on Chapter 4.

**Augusta:**

[https://www.telstarbob.net/bbrown/Math1540\\_daily\\_sch\\_Spring\\_2021AUG.htm](https://www.telstarbob.net/bbrown/Math1540_daily_sch_Spring_2021AUG.htm)

**Swainsboro/Statesboro:**

[https://www.telstarbob.net/bbrown/Math1540\\_daily\\_sch\\_Spring\\_2021SWST.htm](https://www.telstarbob.net/bbrown/Math1540_daily_sch_Spring_2021SWST.htm)

Completed all the homework for Chapter 4.

2. Work the practice test by yourself.


<http://telstarbob.net/bbrown/Math1540OL/CalculusITest3Practice.pdf>

3. Watch Practice test 3 video if you need help on any problem.

<https://www.youtube.com/watch?v=cy92PtEIsYU&feature=youtu.be>

4. Ask me any questions or even request a Zoom Video Tutoring Session if you need additional help.

5. Calculate critical points of a function ( $f' = 0$ , or  $f'$  is not defined)



6. For a function  $y = f(x)$ , determine intervals in which the function is increasing ( $y' > 0$ ) or decreasing ( $y' < 0$ ). Solve  $y' = 0$  to determine the intervals to consider.
7. Be able to calculate any absolute max/min or relative max/min of a function. **Understand that, if the function is defined over a closed interval  $[a, b]$ , you must evaluate the function at the end points as well as the critical points.**
8. Calculate  $x = c$  in the Mean Value Theorem that satisfies  $f'(c) = \frac{f(b)-f(a)}{b-a}$
9. Define intervals on which a function is concave up ( $f'' > 0$ ) or concave down ( $f'' < 0$ ).
10. Use l'Hopital's Rule to find limits. You may have to rewrite so that you have  $\frac{0}{0}$  or  $\frac{\infty}{\infty}$ .
11. Solve max or min applications problems.
12. Calculate antiderivatives.
13. Solve initial value problems given  $y'$  or  $y''$  or position  $s'$  or  $s''$  and determine  $y$  or  $s$  as required.  
You solve by finding the antiderivative, adding a constant each time you integrate, and solve for the constants by using the initial values or values given at a point.

All of these ideas are illustrated in the Practice Test.

On the test, you may have this study guide as well as the derivative and anti-derivative tables at the end of this note.

Good luck,

## Differentiation Rules

$$\frac{d}{dx} x^n = nx^{n-1}$$

$$\frac{d}{dx} \ln x = \frac{1}{x}$$

$$\frac{d}{dx} \sin x = \cos x$$

$$\frac{d}{dx} \cos x = -\sin x$$

$$\frac{d}{dx} \tan x = \sec^2 x$$

$$\frac{d}{dx} \csc x = -\csc x \cot x$$

$$\frac{d}{dx} \sec x = \sec x \tan x$$

$$\frac{d}{dx} \cot x = -\csc^2 x$$

$$\frac{d}{dx} e^x = e^x$$

$$\frac{d}{dx} a^x = (\ln a)a^x$$

$$\frac{d}{dx} \sin^{-1} x = \frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx} \cos^{-1} x = \frac{-1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx} \tan^{-1} x = \frac{1}{1+x^2}$$

$$\frac{d}{dx} \cot^{-1} x = \frac{-1}{1+x^2}$$

$$\frac{d}{dx} \sec^{-1} x = \frac{1}{x\sqrt{x^2-1}}$$

$$\frac{d}{dx} \csc^{-1} x = \frac{-1}{x\sqrt{x^2-1}}$$

Note: **Remember the Chain Rule! If the variable in the function is not just plain x**


**but is u(x), you must multiply the results by  $\frac{du}{dx}$ .**

Product Rule:  $(uv)' = uv' + vu'$

Quotient Rule:  $\left(\frac{u}{v}\right)' = \frac{vu' - uv'}{v^2}$

**TABLE 4.2** Antiderivative formulas,  $k$  a nonzero constant

Function	General antiderivative	Function	General antiderivative
1. $x^n$	$\frac{1}{n+1}x^{n+1} + C, \quad n \neq -1$	8. $e^{kx}$	$\frac{1}{k}e^{kx} + C$
2. $\sin kx$	$-\frac{1}{k}\cos kx + C$	9. $\frac{1}{x}$	$\ln x  + C, \quad x \neq 0$
3. $\cos kx$	$\frac{1}{k}\sin kx + C$	10. $\frac{1}{\sqrt{1-k^2x^2}}$	$\frac{1}{k}\sin^{-1} kx + C$
4. $\sec^2 kx$	$\frac{1}{k}\tan kx + C$	11. $\frac{1}{1+k^2x^2}$	$\frac{1}{k}\tan^{-1} kx + C$
5. $\csc^2 kx$	$-\frac{1}{k}\cot kx + C$	12. $\frac{1}{x\sqrt{k^2x^2-1}}$	$\sec^{-1} kx + C, \quad kx > 1$
6. $\sec kx \tan kx$	$\frac{1}{k}\sec kx + C$	13. $a^{kx}$	$\left(\frac{1}{k \ln a}\right)a^{kx} + C, \quad a > 0, a \neq 1$
7. $\csc kx \cot kx$	$-\frac{1}{k}\csc kx + C$		



**The Unit Circle**